

When Children Won't Eat (and how to help!)



SOS Approach to Feeding Parent & Caregiver Workshop

Presented by Dr. Kay Toomey

Learn more about the SOS Approach to Feeding
at www.sosapproach.com



Resources

Education

Strategies

Support

Bring Fun Back to Mealtimes with the Trusted Feeding Experts

Welcome

- As you take your seat, please pick up a FOOD RANGE handout and the talk packet.
- Please fill out the FOOD RANGE handout as you are waiting for the talk to begin = list in each column (proteins, fruits/veggies, starches) the food your child eats REGULARLY (= 2-3 bites 80% of the time it is served) by recipe or brand name (e.g. chicken piccata or chicken nuggets versus just chicken; Chex cereal versus just cereal; applesauce AND apples versus just apples; garlic bread AND wheat bread versus just bread)

Toomey & Associates, Inc. © 2019

1

WHEN CHILDREN WON'T EAT

(and Dr. Kay Toomeyhelp!)

Copyright protected - 2019
Toomey

Toomey & Associates, Inc. © 2019

2

Statement of Financial & Non-Financial Relationships

Dr. Toomey

- Paid Consultant for Nestec, a division of Nestle (Gerber Child Advocate/Developmental Panel member)
- Unpaid Medical Professional Council member for the Feeding Matters

Lindsay Beckerman

- Board Member for Charlotte's Day

Toomey & Associates, Inc. © 2019

3

Objectives

1. Participants will be able to identify the major reasons why children won't eat.
2. Participants will be able to outline the necessary developmental skills for eating/feeding.
3. Participants will be able to implement at least 3 strategies to improve a child's eating and mealtime participation.

Toomey & Associates, Inc. © 2019

4


When a child doesn't eat well enough to grow or to have good nutrition, it is most often because something about their body is not working quite right.

Dr. Kay Toomey

**IT'S NOT ALL IN THEIR HEAD.
IT'S ALL IN THEIR BODY.**


Toomey & Associates, Inc. © 2019

5



Toomey & Associates, Inc. © 2019

6



7

Development

- Many people say – “it’s just a developmental phase”. “All children are picky and all kids outgrow it”.
 - Not supported by the research
 - ¼ to ½ of children will struggle with some type of feeding difficulty between birth to 10 years of age
 - Only about ½ will “outgrow” their picky eating

Toomey & Associates, Inc. © 2019

8

Development

There are times in development when feeding gets better or worse for most children.

- 4-6 weeks
- 4 - 6 months
- 12 - 14 months
- 18 - 36 months
- 5 - 7 years
- 9 - 11 years

Toomey & Associates, Inc. © 2019

9

Development and Skill

- There are also 5 SPECIFIC skills that children need to learn so a child will eat well.

SKILLS

- ✓ Sensory Tolerance/Exploration
- ✓ Postural Stability
- ✓ Tongue Tip Lateralization
- ✓ Rotary Chewing
- ✓ Positive Mind Set

Toomey & Associates, Inc. © 2019

10

When and Why do children become Picky Eaters or Problem Feeders?

- Children who have identified or unidentified physical issues (even mild), often don't fully manage these developmental transition times, and/or fail to gain the correct skills to eat enough to grow well or have good nutrition.
=> Picky Eating or Problem Feeding

Toomey & Associates, Inc. © 2019

11

Environment

- Parents are NOT the major Cause of children's feeding problems.
 - Research shows Parents causing their child's feeding problems in only 5-10% of the cases.
- Skill Deficits and physical problems are the major causes of feeding difficulties.

Toomey & Associates, Inc. © 2019

12

Environment

THAT SAID ...

- Our Family Mealtime Structure, Routines & Schedules;
- Our Ability to be a Good Role Model;
- The Food Choices we make, and
- How We Talk to our children about food,

ALL play a role in whether our child's feeding issues get better or worse.

Toomey & Associates, Inc. © 2019

13

Physical Issues to consider when a child struggles to eat

- **EATING MYTH**

= "Eating is the body's #1 Priority

- **FACTS:**

Breathing = #1 Priority

- Postural Stability = #2 Priority

Toomey & Associates, Inc. © 2019

14

Organ Systems

- **Respiratory**

- The Body's #1 Priority = Breathing (not eating)
- Immature lungs/oxygen use in the NICU; chronic congestion, colds and/or sinusitis; asthma; suck-swallow-breathe coordination issues

- **Cardiac**

- Heart difficulties can impact oxygenation and may lead to early fatigue during feedings

- **GI (gastrointestinal)**

- If every time you eat it hurts, you learn to not eat.
- If you can't absorb calories, you can't grow.

Toomey & Associates, Inc. © 2019

15

Organ Systems

- Metabolic (ability to use calories ingested)
- Skeletal (impact on postural stability)
- Endocrine (appetite and satiation peptides and hormones)
- Neurological (motor coordination, learning)
- Renal/excretory (ability to process nutrients & wastes)
- Immune System (ability to tolerate the food)

Toomey & Associates, Inc. © 2019

16

Physical Issues to Consider in Feeding

EATING MYTH

= "Eating is the body's #1 Priority"

FACTS:

- Breathing = #1 Priority
- Postural Stability = #2 Priority

Toomey & Associates, Inc. © 2019

17

Children with Muscle Problems (such as Low or Low/Normal Muscle Tone) May...

- Slouch while sitting
- Prop when sitting
- Joint lock/fix
- Slide out from underneath tables/trays
- Not self feed
- Prefer to stand and eat
- Like to walk around and eat
- Appear stronger than they are because of joint locking or fixing

Toomey & Associates, Inc. © 2019

18


Postural Stability Exercise

- Sit up straight
- No arms/hands on table or arm rest of chair
- Pick feet up off the floor and keep them up until instructed to set them back down

Toomey & Associates, Inc. © 2019

19


Postural Stability = a 90-90-90 position



Toomey & Associates, Inc. © 2019

20

90-90-90



Toomey & Associates, Inc. © 2019

21

Seating Recommendations

- Infant Feeding Chair, High Chair, Bouncy Seat or Swing Seat for children age 6 months to 7/8 months (slight tilt back)
- Infant Feeding Chair or High Chair (in upright position) for children 7/8 months to 14/16 months
 - Chicco Polly Progress; Graco Swivi Seat 3-in-1 Booster; Fisher Price SpaceSaver High Chair
- Adjustable Wooden Chair or high chair pushed up to the table for children over 14-16 months
 - Height Right Chair; STOKKE Tripp Trapp; Svan Chair; EURO II Chair

Toomey & Associates, Inc. © 2019

22


Other Postural Stability Strategies

- “No-skid” mat on seat of child’s chair
 - Shelf liner works well
- Foot Rest
 - Cover phone books with duct tape to create a custom foot stool
- Side supports (if needed)
 - Yoga blocks
 - Fill small boxes with packing materials to make them firm, then cover in duct tape

Toomey & Associates, Inc. © 2019

23

Physical Issues to Consider in Feeding: **SENSORY TOLERANCE**



Toomey & Associates, Inc. © 2019

24

SENSORY SYSTEMS: External Environment Information

1. Visual (Vision)
2. Tactile (Touch)
3. Auditory (Hearing)
4. Olfactory (Smell)
5. Gustatory (Taste)

Toomey & Associates, Inc. © 2019

25

SENSORY SYSTEMS: Internal Environment Information


6. Proprioception = position, location, orientation, and movement of the body muscles and joints
7. Vestibular = balance and orientation in space relative to gravity
8. Interoception = ability to read your internal body signals
 - > Sleep
 - > Toileting
 - > Appetite
 - > Temperature recognition
 - > Illness recognition
 - > Emotion regulation

Toomey & Associates, Inc. © 2019

26

STEPS TO EATING

Myth = Eating is a 2 Step Process



Toomey & Associates, Inc. © 2019

27

EATING MYTH = Eating is instinctive

FACT

- Eating is driven primarily by the Appetite instinct only for the first 4-6 weeks of life.
- Eating is then driven by Primitive Motor Reflexes until 4 - 6 months of age.
 - Reflexes for eating “go away” by 6 months
 - Anatomy also changes
- After 6 months of age ...

Eating is a Learned Behavior

Toomey & Associates, Inc. © 2019

28

A child has 3 choices after 4-6 months...

LEARN TO EAT ...

- **Learn to NOT eat**
- **Learn to “kinda, sorta” eat**

29

Because after 4/6 months of age, eating is a learned behavior ...

• **We need to re-think our role as parents/caregivers during meals:**

- Every meal is class
- You are the professor
- Your child is the student
- Food is your subject

Goal = for your child to learn to eat any and all foods over time in order to develop a healthy, lifelong relationship with food

Toomey & Associates, Inc. © 2019

30

Necessary Oral-Motor Skills

- Tongue Tip Lateralization

- Required to transition onto textured table foods
- Required to achieve correct positioning of food onto back molars for chewing

- Rotary Chewing

- Required to eat “real” meat, hard/raw vegetables, and hard raw fruits (especially with peels)

Toomey & Associates, Inc. © 2019

31

Other Critical Developmental Shifts

- 12-14 Months = shift in flavor perception

- 18-36 Months = shift in self awareness

- Aware of self as own person, so really aware of what does and doesn't feel good sensory-wise
- Aware of own opinions and that I can actively do something about it (e.g. saying “no”)
- Cognitive shift from sensori-motor to magical thinking
- Desire to “food jag”

Toomey & Associates, Inc. © 2019

32

Other Critical Developmental Shifts

- 5-7 years = shift to Logical Thinking

- Desire to Food Jag

- 9-11 years = shift to Abstract thought

- Desire to Food Jag

When children shift from one cognitive stage to the next, their sensory functioning regresses. As a result, they want to eat the same exact food over and over because then they don't need to use their sensory brain.

Toomey & Associates, Inc. © 2019

33

Management of Food Jags

- **What are Food Jags?**

When a person eats the same food, prepared the same way, every day or at every meal.

- **The problem with food jags:**

Children eventually get burned out on these foods and they are typically permanently lost out of that child's food range.

Toomey & Associates, Inc. © 2019

34

Preventing Food Jags

- **Offer any one particular food ONLY every OTHER day**

➤ To get through 2 full days without repeating a single food means your child needs 10 different proteins, 10 different starches and 10 different fruits/veggies that they will eat regularly

- So let's look at your food lists

Toomey & Associates, Inc. © 2019

35

Picky Eaters vs Problem Feeders

PICKY EATERS

- Decreased range/variety of foods but will eat ≥ 30 foods
- Foods lost due to "burn out" usually re-gained after 2 weeks
- Tolerates new foods on plate and usually can touch or taste
- Eats ≥ 1 food from most all food texture groups or nutrition groups

PROBLEM FEEDERS

- Restricted range or variety of foods, usually < 20
- Foods lost are NOT re-acquired
- Cries/"falls apart" with new foods
- Refuses entire categories of food textures or nutrition group

Toomey & Associates, Inc. © 2019

36

Picky Eaters vs Problem Feeders

PICKY EATERS

- Adds new foods to repertoire in 15-25 steps
- Typically eats with family, but frequently eats different foods than family
- Sometimes reported as "picky eater" at well child checks

PROBLEM FEEDERS

- Adds new foods in > 25 steps
- Usually eats different foods than family and often eats alone
- Persistently reported as "picky eater" across multiple well child checks

Toomey & Associates, Inc. © 2019

37

General Treatment Strategies

• **Social Modeling:** = Teaching the Social Experience of Eating

1. Family Meals
 - Sit down and eat with your child
 - MEALS ARE CLASSTIME. YOU ARE THE PROFESSOR SO YOU NEED TO MAKE SURE YOU SHOW UP TO CLASS.
2. Model good feeding behaviors

Toomey & Associates, Inc. © 2019

38

Social Modeling

3. Discuss the foods being served
 - REMEMBER – MEALS ARE CLASSTIME, SO YOU NEED TO DISCUSS YOUR SUBJECT
 - Describe how the food looks, feels, smells, tastes, and sounds as you chew it using SCIENCE words
4. Over-exaggerate motor movements
.... AND TEACH YOUR SUBJECT
5. The child is **not** to be the focus
 - THE FOOD IS THE FOCUS OF THIS CLASS, AND LEARNING ABOUT THE FOOD.

Toomey & Associates, Inc. © 2019

39

Social Modeling

6. Imitate your child's eating.

- Otherwise, they are not going to want to come to your class.
- Research shows, children eat new food ONLY when their parent sits down and eats that new food too.

7. Make sure the food is FUN

8. Child should be involved in all aspects of the meal that he/she can

40

Social Modeling

9. Do not punish children at meals

10. Child needs to stay at the table => no "time out" during the meal

WHY? If we stress out the child,
their adrenalin will increase.

41

Issues with Adrenalin

- Causes Appetite suppression
 - Hypothalamus; amygdala; hippocampus; gut all play a role in appetite
- Shuts off digestive tract
- Shifts child into "React mode" versus "learning mode"

42

Structure to Meal and Snack Times

1. Use same place, table and/or designated surface for EACH meal and snack
✓ Buy new placemats as a “cue to eating”
2. Make sure there is at least one very preferred food for your child at EVERY meal and snack (so they have something to easily eat). BUT, also make sure that you offer your child the family foods as well (so they engage in some learning at every meal too).

Toomey & Associates, Inc. © 2019

43

3. Follow same time schedule – a child older than 18 months needs to eat every 2.5 to 3 hours throughout the day

Eating Myth = Children only need to eat 3x's a day

Toomey & Associates, Inc. © 2019

44

WHO Multicentre Growth Reference Study Group, Acta Paediatrica, 2006 (450): 27-37.

“Complementary feeding in the WHO Multicentre Growth Reference Study”

- N = 8440 children from Brazil, Ghana, India, Norway, Oman and USA
- Ages = birth to 24 months
- Mean age for introducing foods = 5.4 months
- Typical # solids meals at 6 months = 2
- Typical # of solids meals at 9 months = 4
- Typical # of solids meals at 12-24 months = 4-5
- 6-12 Months: Average # Total meals per day = 11
 - Total = breast, bottle and solids feedings
- 12-24 Months: Average # Total meals per day = 7

Toomey & Associates, Inc. © 2019

45

Structure to Meal and Snack Times

4. Routine to Family Meals

- Step 1 = warning
- Step 2 = transition
 - = "it is time to wash hands now"
- Step 3 = sit at table with empty plate
- Step 4 = Family Style Serving
 - = everyone gets a little of EVERY food served (use a "Learning Plate if needed)
- Step 5 = Clean Up
 - = throw or blow one piece of every food served at that meal into trash/scraps bowl

Toomey & Associates, Inc. © 2019

46

Structure to Meal and Snack Times

5. Make sure you present at least one Protein food, one food that is a starch and one fruit/vegetable at EVERY meal AND Snack

6. Don't "short order cook"

- Can make 2 meals BUT serve them together as **one** big meal.
- Everyone is expected to take at least one piece of every food served at the table

Toomey & Associates, Inc. © 2019

47

Structure to Meal and Snack Times

7. If your child gets visually overwhelmed easily, present only 3 foods at one time and serve more than one course,

- Make sure that every meal **and** snack has
 - = 1 protein,
 - = 1 starch, AND
 - = 1 fruit/vegetable

Toomey & Associates, Inc. © 2019

48

Structure to Meal and Snack Times

8. Present foods in manageable bites for your child's SKILL level (not chronological age)

- Cut foods into small, soft cubes or long sticks
- Present foods on a cocktail fork
 - Helps get the food onto molars
- Food choppers
 - Helps "pre-chew" foods to decrease rotary chewing, strength and endurance demands

Toomey & Associates, Inc. © 2019

49

Structure to Meal and Snack Times

9. Rule of Thumb = serving size is ONE Tablespoon per year of age (up to the age of 10 years)

- Use a teaspoon measure for calorie dense foods (e.g. peanut butter or cookies, hemp seeds)

10. Limit meals & snacks to 15 to 30 minutes

Toomey & Associates, Inc. © 2019

50

Structure to Meal and Snack Times

11. To prevent Food Jags, do not serve your child the same exact food more than one time each day, and ideally, more than once in 2 days.

- ▶ In order to go two full days without repeating a food, your child needs to have at least 10 proteins, 10 starches, and 10 fruits and vegetables in their food range.

Toomey & Associates, Inc. © 2019

51

RED FLAGS

- does your child need Feeding Therapy

- ☐ Ongoing poor weight gain, weight loss
- ☐ Ongoing choking, gagging, coughing during meals
- ☐ Ongoing problems with vomiting
- ☐ More than once incident of nasal reflux
- ☐ History of a traumatic choking incident

Toomey & Associates, Inc. © 2019

52

RED FLAGS

- ☐ History of eating + breathing problems, with ongoing respiratory issues
- ☐ Inability to transition to baby food purees by 10 months
- ☐ Inability to accept any table food solids by 12 months
- ☐ Inability to transition to a cup by 16 months
- ☐ Has not weaned off most/all baby foods by 16 months

Toomey & Associates, Inc. © 2019

53

RED FLAGS

- ☐ Aversion/avoidance of all foods in specific texture or food group
- ☐ Food range < 20 foods, especially if foods are being dropped
- ☐ Family is fighting about food/feeding
- ☐ Parent repeatedly reports the child as difficult for everyone to feed

Toomey & Associates, Inc. © 2019

54

The SOS Approach to Feeding

Sequential Oral Sensory

And Save Our Ship!

Dr. Kay Toomey

Toomey & Associates, Inc. © 2019

55

TENET 1

= Myths About Eating

Interfere with understanding
and treating feeding problems

Toomey & Associates, Inc. © 2019

56

TENET 2

= Systematic Desensitization

is the best first approach to
feeding treatment

Toomey & Associates, Inc. © 2019

57

Systematic Desensitization

The use of competing “relaxation” responses (= Play with a Purpose) during exposure to a graduated hierarchy of challenging things/stimuli about eating/food (= Steps to Eating) to help a child overcome their difficulties.

Toomey & Associates, Inc. © 2019

58

TENET 3

= the “Normal Development”
of feeding gives us the best
blueprint for feeding treatment

Toomey & Associates, Inc. © 2019

59

TENET 4

= Food Hierarchies/Choices

Play an important role in feeding
treatment

- you can achieve sensory and oral-skill progression with food choices rather than “doing to” the child with stimulation tools
- food hierarchies help sensory systems shift slowly into accepting new foods

Toomey & Associates, Inc. © 2019

60

Please Visit Our Website

www.sosapproach.com

- Access information about SOS trained providers in your area
- Contact the SOS Feeding Team
- Become a Parent/Caregiver Member

Toomey & Associates, Inc. © 2019

Food Range Diary

<u>Child's Name:</u>	<u>Date:</u>
-----------------------------	---------------------

Directions: Please list all of the different foods that your child will eat at least 2-3 bites of, each time that they are served that food (at least ~80 to 90% of the time that it is offered). Please list each individual food by its' name. For example: List chicken nuggets; chicken fingers; chicken lunch meat; chicken pot pie; baked chicken breast VERSUS just listing chicken. List dried or dehydrated fruits or vegetables AND the fresh versions separately. List townie crackers, saltines, graham crackers VERSUS just listing crackers. Don't forget to list what your child drinks. Please be VERY SPECIFIC. Use back of page if extra space is needed.

PROTEINS	STARCHES	FRUITS/VEGETABLES